

Experiments at the International Linear Collider





Daniel Jeans (IPNS/KEK)



ICS2024@APPI

February 2024









Electro-weak transition

Electro-weak physics should contain imprint of physics at much higher energy





Insight through Accelerators.

KEK 2021

Electro-weak transition

Higgs potential changes shape




Higgs particle: excitation of Higgs field

different to all other
fundamental particles
not “matter”, not “force”, no spin




● QUARKS ● LEPTONS ● BOSONS ● HIGGS BOSON



why did the **transition** happen ?

how fast did it happen ?

did it cause the universe's
anti-matter to disappear ?





Is the Higgs potential as we expect

is our **current vacuum**
really stable ?

might the vacuum in our universe
spontaneously decay ?

we've observed all particles of the Standard Model

...but they describe only a small fraction of our universe



To be honest, we understand very little !

a unique window



H_{iggs}
boson

couples to
dark sector ?

a bit different ?



composite
particle ?

one of many
Higgs particles?

exactly as predicted
by today's theory ?


Higgs decay branching fractions

as predicted in the Standard Model




Higgs couples to particles' mass

Deviations in Higgs couplings from BSM physics



new physics @ TeV-scale → few % deviations

Deviations in Higgs couplings from BSM physics



arXiv:1708.08912

→ different BSM models give different deviations


“Higgs Factory”

based on an electron – positron collider

→ **high precision** measurements of Higgs particle and other topics




International Linear Collider



- * electron – positron collider
- * initial energy: 250 GeV (centre of mass)
- * initial luminosity: $1.35 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
- * polarised beams
- * future upgrades in energy and luminosity
~1M Higgs bosons



International Linear Collider



* electron – positron collider

* initial energy: 250 GeV (centre of mass)


* initial luminosity: $1.35 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$

* polarised beams

* future upgrades in energy and luminosity

~1M Higgs bosons

International Linear Collider




* electron – positron collider



why ?
LHC is proton+proton ...

proton - proton



protons are composite: quarks and gluons
→ wide spectrum of q-q , q-g , g-g collision energies

debris from collision of remainder of protons

dominated by “strong force” QCD interactions



e+ e-





elementary particles: each collision has “fixed” energy


almost no “debris”: clean events, easy to analyse

dominated by Electro-Weak interactions






cross-section [fb]




Proton collider

e+ e- collider




International Linear Collider



- * electron – positron collider
- * **initial energy:** 250 GeV (centre of mass)
- * initial luminosity: $1.35 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
- * polarised beams
- * future upgrades in energy and luminosity
~1M Higgs bosons

International Linear Collider






* electron – positron collider

* initial energy: 250 GeV (centre of mass)




why ?

LHC runs at 14 TeV...




what about this process ?



what energy is needed ?


what about this process ?



hint: mass of H boson is 125 GeV

what energy is needed ?


what about this process ?



why is this not so useful ?


hint: Higgs couples to particles' **mass**

International Linear Collider



- * electron – positron collider
- * initial energy: 250 GeV (centre of mass)
- * **initial luminosity: $1.35 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$**
- * polarised beams
- * future upgrades in energy and luminosity
~1M Higgs bosons

International Linear Collider




- * electron – positron collider
- * initial energy: 250 GeV (centre of mass)
- * initial luminosity: $1.35 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$



what is luminosity?

Number of Higgs bosons = cross-section * integrated luminosity
 = cross-section * running time * luminosity

need enough luminosity to get enough Higgs bosons in a reasonable time



350 fb = $350 \times 10^{-15} \times 10^{-24} \text{ cm}^2 = 3.5 \times 10^{-37} \text{ cm}^2$

- ILC luminosity: $1.35 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
- $4.7 \times 10^{-3} \text{ Higgs s}^{-1}$
 - one Higgs every 3~4 minutes
 - 150k per year (if running continuously)

(after a few years,
 plan to upgrade luminosity
 to increase this rate)



N electrons



N positrons



Bunch size

Luminosity ~

$$\frac{N * N * (\text{repetition rate}) * (\text{enhancement factor})}{\text{bunch_size (vertical)} * \text{bunch_size (horizontal)}}$$

Large N

~ 10¹⁰


Large repetition rate

~ 6500 / s

Small bunch size


~ 7 nm (vertical) ~ 500 nm (horizontal)

International Linear Collider



- * electron – positron collider
- * initial energy: 250 GeV (centre of mass)
- * initial luminosity: $1.35 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
- * **polarised beams**
- * future upgrades in energy and luminosity
~1M Higgs bosons

International Linear Collider




- * electron – positron collider
- * initial energy: 250 GeV (centre of mass)
- * initial luminosity: $1.35 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
- * **polarised beams**



what is beam polarisation ?
what's good about it ?

What is beam polarisation?



mostly positive helicity


mostly negative helicity

if highly relativistic:


mostly **right-handed**

mostly **left-handed**


In the electro-weak interactions,
Left and Right-handed fermions are different particles



flipping the beam polarisations → 4 different experiments!



International Linear Collider




- * electron – positron collider
- * initial energy: 250 GeV (centre of mass)
- * initial luminosity: $1.35 \times 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$
- * polarised beams
- * future upgrades in energy and luminosity




why increase the energy ?

Higgs production in electron-positron collisions

associated Higgs production




ZZ & WW fusion





Higgs studies can start at **250 GeV**

full set of Higgs measurements:
add **~500 & ~1000 GeV**


top quark Yukawa



Higgs boson self-coupling



Circular collider (electron-positron)



Energy loss by synchrotron radiation:

$$\text{power loss} \sim E^4 / (m^4 r^2)$$

E: energy

m: particle mass

r: ring radius

- practically limits the maximum beam energy
- difficult to increase energy in a ring
unless what ?

Electrical power $\sim E^4$

Linear Collider



Beam energy limited by tunnel length


- “easy” to extend (reusing existing tunnel)

Electrical power $\sim E$

ILC technology

Key Technology:


Super-Conducting Radio Frequency acceleration



accelerate electrons through 30~35+ million volts every meter

A voltage generator induces an electric field inside the rf cavity. Its voltage oscillates with a radio frequency of 1.3 Gigahertz or 1.3 billion times per second.

The electrons always feel a force in the forward direction.




An electron source injects particles into the cavity in phase with the variable voltage.


The electrons never feel a force in the backward direction.

Super-Conducting cavities for ILC


Super-conductor → dramatically reduce heating
→ more efficient



Niobium :
good superconductor




cryomodule



~900 for ILC


room temperature

keep at 2K (-271°C)



liquid Helium bath

~7,800 for ILC



Superconducting Test Facility (STF)



3 km of “ILC-like” technology
European XFEL @ DESY




cryomodule at FNAL, destined for LCLS-II @ SLAC




e-105

Experiments at ILC

two international groups developing detectors for ILC



design detectors with
unprecedented **precision**
→ enable ILC program



challenging requirements


to maximise physics harvest

- efficiency, identification, resolution
- hadronic jet resolution
- angular coverage


technological advances


- new technologies
- low power, integrated electronics
- compact devices
- machine learning / AI
- quantum sensors





Geant4 simulation in ILD detector






tracking detector

**precise momentum
of *charged* particles**


$d p_T / p_T \sim 3 \times 10^{-5} p_T$



**electromagnetic
calorimeter**

**reasonable precise
measurement of
electrons, positrons,
photons**


$dE/E \sim 20\% / \sqrt{E}$




hadronic calorimeter

**less precise
measurement of
charged and neutral
hadron energies**


$$dE/E \sim 50\% / \sqrt{E}$$




what type
of event ??




??





@ 250 GeV





??



??




$e^+e^- \rightarrow e^+e^- h$ [invisible h decay]
@ 250 GeV





$e^+e^- \rightarrow t\bar{t}h$ [$t\bar{t} \rightarrow 6q$, $h \rightarrow b\bar{b}$]
@ 1000 GeV

Higgs-strahlung process is particularly powerful

- Higgs can be selected by looking **only** at Z decay products
 - we know **initial e^+e^- 4-momentum** (at lepton collider)
 - we precisely measure **4-momentum of Z**
 - we can trivially extract **4-momentum of “H”**
- select Higgs events with **no decay mode bias** (e.g. invisible Higgs)





count total number of produced Higgs events, and extract Higgs mass without looking at Higgs decay products

- not affected by e.g. unexpectedly weird Higgs decays
- “model independent”


many processes will
produce 1 or more
W, Z, H

these usually ($\sim 70\%$)
decay to $q\bar{q}$

→ shower

→ hadronise

→ jets



hadronic jet:

charged hadrons


pions, kaons, protons ...

photons

from π^0 , η , ... decays

neutral hadrons

K^0_L , neutrons, ...



hadronic jet:

charged hadrons

pions, kaons, protons ...

ave. ~65% of energy

photons


from π^0 , eta, ... decays

ave. ~25% of energy

neutral hadrons

K^0_L , neutrons, ...

ave. ~10% of energy



how should we measure jet energy ?

detector performance requirements


track momentum

impact parameter

transparent tracker


jet energy

cover all solid angle around collision




precision on Higgs boson couplings based on realistic simulation and analysis

compared to **HL-LHC-only**



Higgs coupling to ...

model independent fit possible at ILC




arXiv:1903.01629

ILC facility

unique e^- & e^+ beams
high intensity
high energy
high quality

→ potential for studies
beyond Higgs,
beyond particle
physics

new ideas welcome !



Green ILC




linear accelerator and **super-conducting technology** were chosen because they minimize energy loss

none the less, ILC operation requires **111 MW** (at 250 GeV) assuming current energy mix: **320 kton CO₂** per year

- continue development of **energy saving technologies** for ILC
- use of **waste energy** (heat) by local industry
- encourage and prioritize **renewable energy** sources
- encourage local **forestry** industry: wooden construction



proposed Higgs factory projects



ILC project


ILC is a large project

many skilled human resources

extensive production facilities

construction 635.0 – 702.8 GJPY

annual operation 36.6 – 39.2 GJPY



→ a true international project is essential

IUPAP International Union of Pure and Applied Physics



C11 Commission of Particles and Fields



ICFA International Committee for Future Accelerators




ilc International Development Team




P. Campana	Chair	USA
T. Schoerner-Sadenius	Secretary	Germany
P. Sphicas	[ECFA chair]	CERN Member States
F. Gianotti	[CERN DG]	CERN Member States
B. Heinemann		CERN Member States
L. Merminga	[FNAL director]	USA
S. Dasu		USA
N. Roe		USA
I. Koop		Russia
V. Obraztsov		Russia
Y. Wang	[IHEP director]	China
U. Egged		Other Countries
G. Gil da Silveira		Other Countries
T. Nagoya		Japan
M. Yamauchi	[KEK DG]	Japan
R. Teuscher		Canada
F. Canelli	Chair of the IUPAP C-11 (ex officio)	

ILC project promotion in Japan



from late 1980s/'90s → several linear collider studies JLC, GLC, NLC, TESLA, ...





Strong consensus in world-wide HEP community for electron-positron Higgs factory (US, Europe, Japan, ...)

Several Higgs Factory proposals
ILC is most technologically developed option

Japanese government considering whether to host ILC in Japan

candidate ILC site

THE TOHOKU REGION OF JAPAN



selected as candidate site by scientists from Japan and abroad





@Iwate_ilc
@ichinoseki_ilc
@Oshu_ILC
@ILCsupporters
@ilc_tsushin
@LCNewsLine

we want ... **ilc** !!



summary



Higgs particle presents a **once-per-generation opportunity** to look into our universe's beginnings, perhaps its destiny

(most) particle physicists agree it would be great to have a “Higgs Factory”
several such projects under consideration

ILC is an **ideal facility** to enable this study of the Higgs
it requires joint effort from the **worldwide community**:
governments, local communities, industries, academia

ILC uses **technologies** developed around the world
technologies have been proven

Hosting ILC in Iwate/Japan/Asia will promote position at the forefront of
science, technology, culture, and society through the 21st century